

Inoculation experiments with *Puccinia bromina* var. *paucipora* in 1967

Infekční pokusy s *Puccinia bromina* var. *paucipora* v r. 1967

Zdeněk Urban*) and Halvor B. Gjaerum**)

By infectious experiments with aeciospores it was found that *Puccinia bromina* var. *paucipora* Urban forms aecia in Czechoslovakia on *Pulmonaria officinalis* subsp. *obscura* and on *Symphytum tuberosum*, in Norway and Czechoslovakia also on *Symphytum officinale*. As hosts of dikaryotic phase were determined the species *Bromus inermis*, *B. ramosus* subsp. *benekenii* and *B. secalinus*. *Bromus secalinus* as host is known only by experiment.

Infekčními pokusy pomocí aeciospor bylo zjištěno, že *Puccinia bromina* var. *paucipora* Urban vytváří aecia v Československu na *Pulmonaria officinalis* subsp. *obscura* a na *Symphytum tuberosum*, v Norsku a Československu též na *Symphytum officinale*. Jako hostitelé dikaryotické fáze byly stanoveny druhy: *Bromus inermis*, *B. ramosus* subsp. *benekenii* a *B. secalinus*. *Bromus secalinus* je znám jako hostitel jen v pokuse.