

Ophiostoma stenoceras and O. grandicarpum (Ophiostomatales), first records in the Czech Republic

DAVID NOVOTNÝ¹ and PETR ŠRŮTKA²

¹ Research Institute of Crop Production - Division of Plant Medicine, Drnovská 507, 161 06 Praha 6 – Ruzyně, Czech Republic, e-mail: novotny@vurv.cz,

² Department of Forest Protection, Faculty of Forestry, Czech Agricultural University, Kamýcká 129, 165 21 Praha 6 - Suchbátka, Czech Republic

Novotný D. and Šrůtka P. (2004): *Ophiostoma stenoceras* and *O. grandicarpum* (Ophiostomatales), first records in the Czech Republic. - *Czech Mycol.* 56: 19-32

Two species of ophiostomatoid fungi were observed in oaks. *Ophiostoma stenoceras* was isolated during a study of endophytic mycobiota of the roots and seedlings of a sessile oak (*Quercus petraea*). *Ophiostoma grandicarpum* was recorded in the stem of a pedunculate oak (*Q. robur*). These fungi have not yet been reported from the Czech Republic. The knowledge on the occurrence of ophiostomatoid fungi in the Czech Republic is reviewed.

Key words: ophiostomatoid fungi, distribution, oak, roots, bark, *Ceratocystis*, *Quercus petraea*, *Quercus robur*